

SYMPTOMS of Strangulation

Source: Strangulation in Intimate Partner Violence, Chapter 16, Intimate Partner Violence, Oxford University Press, Inc. 2009.


Injuries not visible to the naked eye; may be observable only to the victim.*



strangulationtraininginstitute.com

NEUROLOGICAL

- Loss of memory
- Loss of consciousness
- Behavioral changes
- Loss of sensation
- Extremity weakness
- Difficulty speaking
- Fainting
- Urination
- Defecation
- Vomiting
- Dizziness
- Headaches



VOICE & THROAT CHANGE

- Raspy or hoarse voice
- Unable to speak
- Trouble swallowing
- Painful to swallow
- Clearing the throat
- Coughing
- Nausea
- Drooling
- Sore throat
- Stridor

BREATHING CHANGES

- Difficulty breathing
- Respiratory distress
- Unable to breathe

* Visible signs may also be present.

Original artwork and design by Yesenia Aceves

This project is supported all or in part by Grant No. 2016-TA-AX-K067 awarded by the Office on Violence Against Women, U.S. Department of Justice. The opinions, findings, conclusions, and recommendations expressed in this publication/program/exhibition are those of the author(s) and do not necessarily reflect the views of the Department of Justice, Office on Violence Against Women.